arXiv:1104.0401v2 [math.CO] 16 Apr 2011

The second Voronoi conjecture on parallelohedra for zonotopes.[*]

A. Garber[†]
e-mail: alexeygarber@gmail.com

January 18, 2013

Abstract

We prove the second Voronoi conjecture on parallelohedra for zonotope. We show that for a given face-to-face tiling of d-dimensional Euclidean space into parallel copies of zonotope Z there are d vectors, connecting centers of zonotopes with common facet, that are basis of the correspondent lattice of the tiling.

AMS subject classification: 52B20, 52C22.

1 Parallelohedra

Definition 1.1. A polytope $P \subseteq \mathbb{R}^d$ is called a *parallelohedron* if Euclidean space $\mathbb{R}^d$ can be tiled into non-overlapping parallel copies of P.

There are several classical results in the theory of parallelohedra. Here we will mention two of them. The first one is the Minkowski theorem [4] that claims that parallelohedron P is centrally symmetric, any facet of P is centrally symmetric and projection of P along any of its ridge (i.e. face of codimension 2) is either parallelogram or centrally symmetric hexagon. The second one is the Venkov theorem that claims that these three conditions of Minkowski are sufficient for P to be a parallelohedron.

In general case two parallelohedra with common boundary point can share only a part of faces of both, as in usual brickwork two bricks from consecutive horizontal layers share only half of facet.

Definition 1.2. If intersection of any two polytopes of the tiling $\mathcal{T}$ of $\mathbb{R}^d$ into non-overlapping polytopes is a face of both of them (this face can be empty) then the tiling $\mathcal{T}$ is called *face-to-face* tiling.

---

[*] This work is financially supported by RFBR (projects 11-01-00633-a and 11-01-00735-a) and by grant of the President of the Russian Federation НШ-5413.2010.1.

[†] Lomonosov Moscow State University, Faculty of Mechanics and Mathematics, Department of Higher Geometry and Topology. Russia, 119991, Moscow, Vorob'evy gory, 1, A-1620.

McMullen showed [3] that if parallelohedron P admits an arbitrary tiling of $\mathbb{R}^d$ into parallel copies then it admits also and a face-to-face tilings, namely McMullen proved that Minkowski conditions are necessary for parallelohedron P with non face-to-face tiling. Further for a given parallelohedron P we will consider only correspondent face-to-face tiling $\mathcal{T}(P)$. The uniqueness (up to translation) of such face-to-face tiling into copies of P is evident. Centers of all polytopes of the tiling $\mathcal{T}(P)$ forms a d-dimensional lattice $\Lambda(P)$.

Also for an arbitrary d-dimensional lattice Λ we can construct a d-dimensional parallelohedron. For a fixed point $O \in \Lambda$ consider a polytope P_Λ that consists of all points of $\mathbb{R}^d$ that are closer to O than to any other point of Λ. The constructed polytope DV_Λ is called the *Dirichlet-Voronoi polytope* for Λ and this polytope is a parallelohedron because for different lattice points corespondent polytopes differs only by translation and all such polytopes gives us a face-to-face tiling of $\mathbb{R}^d$.

The first conjecture of Voronoi states that all parallelohedra can be obtained from Dirichlet-Voronoi polytopes with affine transformations.

Conjecture 1 (G.Voronoi [5]). *For any parallelohedron P there exists a lattice Λ and an affine transformation $\mathcal{A}$ such that the polytope $\mathcal{A}(P)$ is the Dirichlet-Voronoi polytope for Λ.*

Consider the set $\mathcal{N}(P)$ of vectors connecting the center of a given parallelohedron P with centers of all other parallelohedra of tiling $\mathcal{T}(P)$ that shares facets with P. It is clear that $\mathcal{N}(P)$ generates the lattice $\Lambda(P)$ because any vector from $\Lambda(P)$ can be represented as a sum of several vectors from $\mathcal{N}(P)$.

Conjecture 2 (G.Voronoi [5]). *We can choose d vectors from $\mathcal{N}(P)$ that forms a basis of $\Lambda(P)$.*

In this paper we will prove the second conjecture of Voronoi in the case of space-filling zonotopes.

2 Delone tilings and Dirichlet-Voronoi tilings

We can generalize the construction of Dirichlet-Voronoi polytope on the case of general positive definite quadratic form.

Definition 2.1. Let $\varphi : \mathbb{R}^d \longrightarrow \mathbb{R}$ be a positive definite quadratic form in d-dimensional Euclidean space and let O be a point of some fixed lattice Λ. The *Dirichlet-Voronoi polytope* $P_\varphi(\Lambda)$ for Λ with respect to form φ is a polytope that consist of all points X of $\mathbb{R}^d$ such that value of φ on vector $\overrightarrow{OX}$ is not greater than value $\varphi(\overrightarrow{O'X})$ for any other point $O' \in \Lambda$.

The constructed polytope $P_\varphi(\Lambda)$ is a parallelohedron too and the correspondent tiling is called a *Dirichlet-Voronoi tiling* $\mathcal{V}_\varphi(\Lambda)$. If we will take φ equals to the usual metric form $\varphi(\mathbf{x}) = \mathbf{x}^T\mathbf{x}$ then we will get the usual Dirichlet-Voronoi polytope DV_Λ.

Applying an affine transformation $\mathcal{A}$ to a given lattice Λ and a polytope $P_\varphi(\Lambda)$ with form $\varphi(\mathbf{x}) = \mathbf{x}^TQ\mathbf{x}$ we will obtain lattice $\mathcal{A}\Lambda$ and polytope $P_{\varphi_\mathcal{A}}(\mathcal{A}\Lambda)$ with respect to quadratic form $\varphi_\mathcal{A}(\mathbf{x}) = (\mathcal{A}^{-1}\mathbf{x})^TQ(\mathcal{A}^{-1}\mathbf{x})$. So to prove the first conjecture of Voronoi for a given polytope P with lattice Λ it is enough to show that there exist a quadratic form φ such that $P = P_\varphi(\Lambda)$.

For any Dirichlet-Voronoi tiling of $\mathbb{R}^d$ into polytopes $P_\varphi(\Lambda)$ we can construct a dual Delone tiling. The Delone tiling $\mathcal{D}_\varphi(\Lambda)$ is defined by lattice Λ and positive definite quadratic form φ. For a given form φ consider an ellipsoid $\mathcal{E}_\varphi$ with equation $\varphi(\mathbf{x}) = 1$. Consider an arbitrary homothetic copy $\mathcal{E}_0$ of $\mathcal{E}_\varphi$ such that $\mathcal{E}_0$ has some d-dimensional set of points from Λ on the boundary $\partial\mathcal{E}$ but does not have points from Λ inside $\mathcal{E}_0$. This ellipsoid $\mathcal{E}_0$ defines a convex polytope inscribed in $\mathcal{E}$ with vertices from the lattice Λ. The set of all such polytopes inscribed in "empty" ellipsoids we will call the *Delone tiling* $\mathcal{D}_\varphi(\Lambda)$.

For any vertex O of the Dirichlet-Voronoi tiling values of quadratic form φ on vectors that connects O with centers of polytopes from $\mathcal{V}_\varphi(\Lambda)$ that meets at O are equal so O will be center of empty ellipsoid that defines one polytope from the Delone tiling $\mathcal{D}_\varphi(\Lambda)$.

3 Zonotopes and dicings

Definition 3.1. A polytope $P \subseteq \mathbb{R}^d$ is called a *zonotope* if P can be represented as a Minkowski sum of finite number of segments. The Minkowski sum of segments defined by vectors of the set $V = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ we will denote by $Z(V)$. Vectors $\mathbf{v}_i$ is called the *zone vectors* of the zonotope $Z(V)$.

Equivalently zonotope can be defined as a projection of cube C^n of some dimension $n \geq d$.

Erdahl in his work [2] proved the first conjecture of Voronoi for zonotopes. Later Deza and Grishukhin proved the first conjecture of Voronoi for zonotopes in terms of oriented matroids [1]. In this paper we will formulate and use several notions and theorems concerning zonotopes and correspondent Delone tilings and Dirichlet-Voronoi tilings from Erdahl's work [2].

Definition 3.2. Consider n families of hyperplanes in d-dimensional Euclidean space. Assume that every family consists of parallel hyperplanes and slices $\mathbb{R}^d$ into layers of a constant width (this width can vary for different families). This set of n families and the correspondent tiling of $\mathbb{R}^d$ is called *dicing* if there are d linearly independent normals to these hyperplanes and every point that belongs to hyperplanes of d families with independent normals also belongs to hyperplanes of all other families.

Assume that the point with radius vector $\mathbf{a}_0$ belongs to hyperplanes of all families then for a given dicing we can choose normals $\mathbf{d}_1, \dots, \mathbf{d}_n$ to families in such a way that all hyperplanes of dicing will be defined by equations $\mathbf{d}_i \cdot (\mathbf{x} - \mathbf{a}_0) = a$ for various integer a. Also we can substitute any vector $\mathbf{d}_i$ by its negative $-\mathbf{d}_i$.

Definition 3.3. In that case we will denote our dicing by $\mathfrak{D}(\mathbf{d}_1, \dots, \mathbf{d}_n)$ and the set $\mathcal{D} = \{\pm\mathbf{d}_1, \dots, \pm\mathbf{d}_n\}$ we will call the *set of normals* of the dicing $\mathfrak{D}$.

Consider an arbitrary vertex of a dicing $\mathfrak{D}$ (i.e. point of intersection of hyperplanes from d families with independent normals) and consider all edges of $\mathfrak{D}$ that incident to this vertex. This set of edges will be centrally symmetric.

Definition 3.4. The mentioned set of edges $\mathcal{E} = \{\pm\mathbf{e}_1, \dots, \pm\mathbf{e}_k\}$ is called the *edge set* of a dicing $\mathfrak{D}$. It is clear that construction of the set $\mathcal{E}$ does not depend on the vertex of a dicing.

Also this construction immediately follows that all vertices of a dicing $\mathfrak{D}$ forms a lattice $\Lambda(\mathfrak{D})$ that we will call the *lattice of a dicing.*

In particulary Erdahl prooved the following theorem on connection between sets $\mathcal{D}$ and $\mathcal{E}$ [2, Theorem 3.1].

Theorem 3.5 (R. Erdahl)**.** *Given set $\mathcal{D}$ of n vectors with d linearly independent vectors can be a set of normals for a dicing if and only if there exists a set $\mathcal{E}$ such that:*

(E1) *Any pair of opposite vectors $\pm\mathbf{e}_i \in \mathcal{E}$ lies in some one-dimensional intersection $\mathbf{d}_{i_1}^{\perp} \cap \ldots \cap \mathbf{d}_{i_{d-1}}^{\perp}$ with independent vectors $d_{i_j} \in \mathcal{D}$ and conversely for any $d-1$ linearly independent vectors from $\mathcal{D}$ there is a correspondent pair of opposite vectors from $\mathcal{E}$;*

(E2) *For any pair of vectors $\mathbf{d} \in \mathcal{D}$ and $\mathbf{e} \in \mathcal{E}$ the scalar product $\mathbf{d}^T\mathbf{e}$ equals to 0 or ± 1.*

Definition 3.6. Matrices $\mathbf{D}$ and $\mathbf{E}$ with vectors from $\mathcal{D}$ and $\mathcal{E}$ written in columns are called *matrix of normal vectors* and *matrix of edge vectors* of the dicing $\mathfrak{D}$ respectively.

Under the affine transformation of $\mathbb{R}^d$ with matrix L matrices $\mathbf{D}$ and $\mathbf{E}$ changes into matrices $\mathbf{D}' = (L^{-1})^T\mathbf{D}$ and $\mathbf{E}' = L\mathbf{E}$. Moreover, there exists an affine transformations such that entries of matrices $\mathbf{D}$ and $\mathbf{E}$ after this transformation will be only 0 and ± 1 [2, Theorem 3.3].

Theorem 3.7 (R. Erdahl)**.** *There exists an affine transformation that will give us a totally unimodular matrix $\mathcal{D}'$, i.e. any of its minor will be equal to 0 or ± 1. Moreover this transformation can be chosen in such a way that both sets $\mathbf{D}$ and $\mathbf{E}$ will contain d vectors of standard basis of $\mathbb{R}^d$, i.e. vectors $(1, 0, \ldots, 0)^T, (0, 1, 0, \ldots, 0)^T, \ldots, (0, \ldots, 0, 1)^T$. Also after this transformation all entries of matrices $\mathbf{D}$ and $\mathbf{E}$ will become 0 or ± 1.*

We will take only a half of vectors from the set $\mathcal{D}$ with no opposite vectors included. This new set $\mathcal{D}^+$ determines the same unique dicing $\mathfrak{D}$. Consider a quadratic form

$$\varphi(\mathbf{x}) = \sum_{\mathbf{d} \in \mathcal{D}^+} \omega_{\mathbf{d}} (\mathbf{d} \cdot \mathbf{x})^2$$

for some positive constants $\omega_{\mathbf{d}}$. The correspondent Dirichlet-Voronoi polytope described in [2, Theorem 4.3].

Theorem 3.8 (R. Erdahl)**.** *The Dirichlet-Voronoi polytope for lattice $\Lambda(\mathfrak{D})$ with respect to quadratic form $\varphi(\mathbf{x})$ is a zonotope with zone vectors*

$$\mathbf{z}_{\mathbf{d}} = \left(\sum_{\mathbf{d} \in \mathcal{D}^+} \omega_{\mathbf{d}} \mathbf{d}\mathbf{d}^T \right)^{-1} \omega_{\mathbf{d}} \mathbf{d}, \quad \mathbf{d} \in \mathcal{D}^+.$$

In the same work Erdahl proved that if $\varphi(\mathbf{x})$ is the standard Euclidean metrics then the zone vectors of the correspondent zonotope are $\mathbf{z}_{\mathbf{d}} = \omega_{\mathbf{d}} \mathbf{d}$ ([2, Sect. 6, p. 442]).

Also [2, Theorem 1.2] claims that Dirichlet-Voronoi polytope for a lattice is a zonotope if and only if the correspondent Delone tiling is a dicing.

4 The second conjecture of Voronoi for zonotopes

Lemma 4.1. *Let $Z = Z(V)$ be a d-dimensional zonotope. Any facet of Z is generated by some $(d-1)$-dimensional subset U of V and conversely any $(d-1)$-dimensional subset of V generates facet $Z(U)$ of the zonotope $Z(V)$.*

Proof. Let π be a hyperplane of some facet of Z. The zonotope $Z(V) = Z(\mathbf{v}_1, \ldots, \mathbf{v}_n)$ is a projection of a cube $C^n \subset \mathbb{R}^n$ onto space $\mathbb{R}^d$ along $(n-d)$-dimensional subspace ψ. Consider the hyperplane $\pi \times \psi$ in the space $\mathbb{R}^n$. This hyperplane is a supporting plane of the cube C^n and hence it defines its face F. The face F is a cube of some dimension and this face is generated by edges of C^n that projects onto vectors of the set V that are parallel to π. Hence the face F is projected into parallel copy of a zonotope $Z(U)$ for some $(d-1)$-dimensional subset U of V. The converse statement can be proven in the analogous way. ☐

Theorem 4.2. *The conjecture 2 is true for space filling zonotopes.*

Proof. Assume that Z is a d-dimensional space filling zonotope, i.e. zonotope that is also a parallelohedron. Then the first conjecture of Voronoi is true for Z [2, Theorem 1.1], i.e. there exists an affine transformation $\mathcal{A}$ such that the zonotope $\mathcal{A}Z$ is Dirichlet-Voronoi polytope of some lattice Λ with respect to the usual Euclidean metrics as a quadratic form. The Delone tiling for Λ with respect to Euclidean metrics is a dicing $\mathfrak{D} = \mathcal{D}(\pm\mathbf{d}_1, \ldots, \pm\mathbf{d}_n)$ so due to theorem 3.8 the zone vectors of the zonotope $\mathcal{A}Z$ can be written as $\omega_i\mathbf{d}_i$.

Any vector that connects centers O_1 and O_2 of copies of $\mathcal{A}Z$ with a joint facet F is perpendicular to this facet because any point of F is equidistant from O_1 and O_2. By lemma 4.1 there are $d-1$ vectors from the set $\{\omega_i\mathbf{d}_i\}_{i=1}^n$ that are parallel to F so by theorem 3.5 there is a vector from $\mathcal{E}$ perpendicular to F. That means the the vector $\overrightarrow{O_1O_2}$ lies in $\mathcal{E}$. The converse statement is also true, if we take any vector $\mathbf{x}$ from $\mathcal{E}$ then $d-1$ linearly independent vectors from $\mathcal{D}$ that are perpendicular to $\mathbf{x}$ will determine a facet of Z. Therefore sets $\mathcal{E}$ and $\mathcal{N}(Z)$ coincides.

Consider a unimodular representation of the dicing $\mathfrak{D}$. In this representation the set $\mathcal{E}$ contains the standard basis of the space $\mathbb{R}^d$ and all other vectors of $\mathcal{E}$ has integer coordinates. Hence in unimodular representation the set $\mathcal{N}(Z)$ contains a standard basis of $\mathbb{R}^d$ and all other vectors of this set has integer coordinates in this basis. So in unimodular representation the lattice $\Lambda(Z)$ coincides with $\mathbb{Z}^d$ and $\mathcal{N}(Z)$ contains its basis. The last property is invariant under linear transformations and changing of the basis in $\mathbb{R}^d$ and this proves the theorem. ☐

References

[1] M. Deza, V. Grishukhin, *Once more about Voronoi's conjecture and space tiling zonotopes.* e-print, http://arxiv.org/abs/math/0203124, 2002.

[2] R. Erdahl, *Zonotopes, Dicings, and Voronoi's Conjecture on Parallelohedra*, Eur. J. of Comb., Vol. 20, N. 6, 1999, pp. 527-549.

[3] P.McMullen, *Convex bodies which tile space by translation.* Mathematika, vol. 27, n. 1, 1980, pp. 113-121.

[4] H. Minkowski, *Allgemeine Lehrsätze über die convexen Polyeder.* Gött. Nachr., 1897, pp. 198-219.

[5] G. Voronoi, *Nouvelles applications des paramètres continus à la théorie des formes quadratiques. Deuxième mémoire. Recherches sur les paralléloèdres primitifs.* J. für Math., vol. 136, 1909, pp. 67-178.

[6] B.A. Venkov, *About one class of Euclidean polytopes* (in Russian). Vestnik Leningr. Univ., ser. Math., Phys., Chem., 1954, vol. 9, pp. 11-31.

arXiv:1104.0401v2 [math.CO] 16 Apr 2011

Вторая гипотеза Вороного о параллелоэдрах для зонотопов.*

А.И. Гарбер†
`e-mail: alexeygarber@gmail.com`

18 января 2013 г.

Аннотация

В данной работе доказана вторая гипотеза Вороного о параллелоэдрах для зонотопов, то есть доказано, что для данного разбиения "грань-в-грань" d-мерного пространства на копии зонотопа Z можно выбрать d векторов, соединяющие центры зонотопов соседних по гиперграням, которые являются базисом соответствующей решетки разбиения.

1 Параллелоэдры.

Определение 1.1. Многогранник $P \subseteq \mathbb{R}^d$ называется *параллелоэдром* если пространство $\mathbb{R}^d$ можно разбить на параллельные копии P, не пересекающиеся по внутренним точкам.

Классическими результатами в теории параллелоэдров являются теорема Минковского [4], которая утверждает, что если многогранник P является параллелоэдром, то он сам имеет центр симметрии, любая его гипергрань имеет центр симметрии и ортогональная проекция P вдоль любой грани коразмерности 2 есть параллелограмм или центрально-симметричный шестиугольник, и теорема Венкова [6], которая утверждает, что данные три условия являются достаточными для того, чтобы многогранник P был параллелоэдром.

В общем случае два соседних параллелоэдра могут пересекаться по части грани каждого из них, например, в обычной "кирпичной кладке" два соседних кирпича из разных горизонтальных слоев пересекаются не по полной грани.

Определение 1.2. Если в разбиении $\mathcal{T}$ любые два параллелоэдра пересекаются по грани каждого из них (эта грань может быть и пустой), то соответствующее $\mathcal{T}$ называется разбиением *грань-в-грань*.

---

[*] Работа выполнена при финансовой поддержке РФФИ (проекты 11-01-00633-а и 11-01-00735-а), а также при поддержке гранта Президента РФ НШ-5413.2010.1.

[†] Московский государственный университет им. М.В.Ломоносова, механико-математический факультет, каф. высшей геометрии и топологии. Россия, 119991, Москва, Воробьевы горы, 1, А-1620.

МакМаллен [3] доказал, что если данный параллелоэдр P допускает разбиение пространства $\mathbb{R}^d$ на параллельные копии произвольным образом, то он допускает и разбиение грань-в-грань, а именно МакМаллен доказал, что даже для параллелоэдра P и разбиения не грань-в-грань три условия Минковского являются необходимыми. В дальнейшем для каждого параллелоэдра P мы будем рассматривать только соответствующее ему разбиение $\mathcal{T}(P)$ грань-в-грань. Единственность такого разбиения очевидна. При этом центры многогранников разбиения $\mathcal{T}(P)$ образуют d-мерную решетку $\Lambda(P)$.

Также по любой d-мерной решетке Λ можно построить d-мерный параллелоэдр. Для фиксированной точки $O \in \Lambda$ рассмотрим многогранник P_Λ, состоящий из всех точек пространства $\mathbb{R}^d$, которые ближе к O чем к остальным точкам Λ. Данный многогранник называется *многогранником Дирихле-Вороного* и является параллелоэдром, так как многогранники для различных точек решетки отличаются параллельным переносом и дают разбиение “грань-в-грань” пространства $\mathbb{R}^d$.

Первая гипотеза Вороного утверждает, что все параллелоэдры могут быть получены с помощью аффинных преобразований из многогранников Дирихле-Вороного.

Гипотеза 1 (Г.Вороной [5]). *Для любого параллелоэдра P существуют такие решетка Λ и аффинное преобразование $\mathcal{A}$, что многогранник $\mathcal{A}(P)$ является многогранником Дирихле-Вороного для решетки Λ.*

Пусть $\mathcal{N}(P)$ — набор векторов соединяющих центр данного параллелоэдра P с центрами его соседями по гиперграням в разбиении $\mathcal{T}(P)$. Очевидно, что множество $\mathcal{N}(P)$ порождает решетку $\Lambda(P)$ так как любой вектор, соединяющий центры двух параллелоэдров разбиения $\mathcal{T}(P)$, можно представить в виде суммы нескольких векторов из $\mathcal{N}(P)$.

Гипотеза 2 (Г.Вороной [5]). *В множестве $\mathcal{N}(P)$ можно выбрать d векторов, являющиеся базисом решетки Λ.*

В данной работе мы докажем вторую гипотезу Вороного в частном случае зонотопов, являющихся параллелоэдрами.

2 Разбиения Делоне и Вороного

Построение многогранника Дирихле-Вороного из предыдущего раздела можно обобщить на случай произвольной квадратичной формы.

Определение 2.1. Пусть $\varphi : \mathbb{R}^d \longrightarrow \mathbb{R}$ — положительно определенная квадратичная форма в $\mathbb{R}^d$ и O — некоторая точка фиксированной решетки Λ. *Многогранником Дирихле* $P_\varphi(\Lambda)$ для данной решетки Λ относительно формы φ называется многогранник, состоящий из всех точек X пространства $\mathbb{R}^d$, для которых значение формы φ на векторе $\overrightarrow{OX}$ не больше чем значение $\varphi(\overrightarrow{O'X})$ для любой точки $O' \in \Lambda$.

Построенный многогранник $P_\varphi(\Lambda)$ также является параллелоэдром и соответствующее разбиение называется *разбиением Дирихле-Вороного* $\mathcal{V}_\varphi(\Lambda)$. Обычный многогранник

Дирихле-Вороного получится если в качестве формы φ рассмотреть обычную форму метрики $\varphi(\mathbf{x}) = \mathbf{x}^T\mathbf{x}$. При этом, если к данной решетке и данному многограннику $P_\varphi(\Lambda)$ с формой $\varphi(\mathbf{x}) = \mathbf{x}^TQ\mathbf{x}$ мы применим аффинное преобразование $\mathcal{A}$, то в результате получится решетка $\mathcal{A}\Lambda$ и многогранник $P_{\mathcal{A}^{-1}\varphi}(\mathcal{A}\Lambda)$, где форма $\mathcal{A}^{-1}\varphi$ задается равенством $\mathcal{A}^{-1}\varphi(\mathbf{x}) = (\mathcal{A}^{-1}\mathbf{x})^TQ(\mathcal{A}^{-1}\mathbf{x})$. Следовательно, для того, чтобы доказать первую гипотезу Вороного для данного параллелоэдра P с данной решеткой Λ достаточно найти такую квадратичную форму φ, что $P = P_\varphi(\Lambda)$.

Двойственным к разбиению $\mathcal{V}_\varphi(\Lambda)$ на многогранники Дирихле-Вороного $P_\varphi(\Lambda)$ является разбиение Делоне, также определяемое по данным решетке Λ и положительно определенной квадратичной форме φ. Квадратичная форма φ определяет эллипсоид $\mathcal{E}_\varphi$, задаваемый уравнением $\varphi(\mathbf{x}) = 1$. Рассмотрим какой-нибудь эллипсоид гомотетичный $\mathcal{E}_\varphi$ содержащий на границе d-мерное подмножество точек из Λ, но не содержащий внутри себя ни одной точки Λ. Такой эллипсоид определяет вписанный в него выпуклый многогранник. Множество таких многогранников, вписанных в "пустые" эллипсоид называется *разбиением Делоне* $\mathcal{D}_\varphi(\Lambda)$.

Если мы рассмотрим произвольную вершину O разбиения Дирихле-Вороного, то значение квадратичной формы φ на векторах, соединяющих O с центрами многогранников из $\mathcal{V}_\varphi(\Lambda)$ ее содержащих, равны, а значит O и будет центром "пустого" эллипсоида, определяющего один из многогранников разбиения Делоне $\mathcal{D}_\varphi(\Lambda)$.

3 Зонотопы и дайсинги

Определение 3.1. Многогранник $P \subseteq \mathbb{R}^d$ называется *зонотопом* если его можно представить в виде суммы Минковского некоторого конечного набора отрезков. Зонотоп, являющийся суммой Минковского отрезков, образованных векторами множества $V = \{\mathbf{v}_1, \ldots, \mathbf{v}_n\}$, обозначается $Z(V)$. Векторы $\mathbf{v}_i$ называются *векторами зон.*

Эквивалентным образом зонотоп можно определить как проекцию куба C^n некоторой размерности $n \geq d$.

Эрдал в работе [2] доказал первую гипотезу Вороного для зонотопов, являющихся параллелоэдрами. Позднее доказательство первой гипотезы Вороного для зонотопов было дано Деза и Гришухиным в терминах ориентированных матроидов [1]. В данной работе мы сформулируем и будем использовать некоторые факты о зонотопах и соответствующих разбиений Делоне и Дирихле-Вороного из указанной работы [2].

Определение 3.2. Рассмотрим в d-мерном пространстве n семейств гиперплоскостей с нормалями $\mathbf{d}_1, \ldots, \mathbf{d}_n$ причем в каждом семействе все плоскости параллельны и дают разбиение пространства на слои равной толщины. Данный набор семейств и разбиение, которое они задают называются *дайсингом* $\mathfrak{D}(\mathbf{d}_1, \ldots, \mathbf{d}_n)$ если среди данных семейств можно выбрать d с линейно независимыми нормалями и через любую точку, через которую проходят плоскости d семейств, проходят плоскости и остальных семейств.

Пусть точка с радиус-вектором $\mathbf{a}_0$ лежит на гиперплоскостях всех семейств дайсинга, тогда для данного дайсинга можно выбрать нормали $\mathbf{d}_1, \ldots, \mathbf{d}_n$ к гиперплоскостям его семейств таки образом, что все гиперплоскости дайсинга задаются уравнениями $\mathbf{d}_i \cdot (\mathbf{x} - \mathbf{a}_0) =$

a для всевозможных целых параметров a. При этом каждый вектор $\mathbf{d}_i$ можно заменить на ему противоположный $-\mathbf{d}_i$.

Определение 3.3. В этом случае множество векторов $\mathcal{D} = \{\pm\mathbf{d}_1, \ldots, \pm\mathbf{d}_n\}$ называется *множеством нормалей* дайсинга $\mathfrak{D}(\mathbf{d}_1, \ldots, \mathbf{d}_n)$.

Выберем произвольную вершину дайсинга $\mathfrak{D}$ (то есть точку пересечения гиперплоскостей из d семейств) и рассмотрим все ребра дайсинга, которые из нее выходят. Это множество ребер будет центрально симметричным.

Определение 3.4. Данное множество $\mathcal{E} = \{\pm\mathbf{e}_1, \ldots, \pm\mathbf{e}_k\}$ называется *множеством ребер* дайсинга. Очевидно, что построение этого множества не зависит от того какую мы вершину выбрали изначально.

Также отсюда следует, что все вершины дайсинга $\mathfrak{D}$ образуют решетку $\Lambda(\mathfrak{D})$, которая называется *решеткой дайсинга.*

В частности Эрдал доказал следующую теорему о связи множеств $\mathcal{D}$ и $\mathcal{E}$ [2, Теорема 3.1].

Теорема 3.5 (Р. Эрдал)**.** *Множество $\mathcal{D}$ с d линейно независимыми векторами является множеством нормалей некоторого дайсинга если и только если существует такое множество $\mathcal{E}$, что:*

(E1) *Любая пара противоположных векторов $\pm\mathbf{e}_i \in \mathcal{E}$ принадлежит одномерному пересечению $\mathbf{d}_{i_1}^{\perp} \cap \ldots \cap \mathbf{d}_{i_{d-1}}^{\perp}$, где векторы $d_{i_j} \in \mathcal{D}$ и линейно независимы и наоборот для любых $d-1$ независимого вектора из $\mathcal{D}$ найдется соответствующая пара противоположных векторов из $\mathcal{E}$;*

(E2) *Для любых векторов $\mathbf{d} \in \mathcal{D}$ и $\mathbf{e} \in \mathcal{E}$ верно равенство $\mathbf{d} \cdot \mathbf{e} = 0, \pm 1$.*

Определение 3.6. Матрицы $\mathbf{D}$ и $\mathbf{E}$, столбцы которых состоят из векторов множеств $\mathcal{D}$ и $\mathcal{E}$ соответственно, называются матрицами *векторов нормалей* и *векторов ребер* дайсинга $\mathfrak{D}$.

При аффинном преобразовании пространства $\mathbb{R}^d$ с матрицей L матрицы $\mathbf{D}$ и $\mathbf{E}$ меняются по правилам $\mathbf{D}' = (L^{-1})^T\mathbf{D}$ и $\mathbf{E}' = L\mathbf{E}$. Более того, существует аффинное преобразование после которого матрицы $\mathbf{D}$ и $\mathbf{E}$ состоят только из 0 и ± 1 [2, Теорема 3.3].

Теорема 3.7 (Р. Эрдал)**.** *Существует аффинное преобразование, после которого матрица $\mathbf{D}'$ будет полностью унимодулярной, то есть любой ее минор будет равен* 0 *или* ± 1. *Более того, это преобразование можно выбрать таким образом, что в каждом из множеств $\mathcal{D}$ и $\mathcal{E}$ можно выбрать по d векторов, которые являются стандартным базисом пространства $\mathbb{R}^d$, то есть векторами $(1, 0, \ldots, 0)^T, (0, 1, 0, \ldots, 0)^T, \ldots, (0, \ldots, 0, 1)^T$. Также в этом базисе элементами матриц $\mathbf{D}$ и $\mathbf{E}$ будут* 0 *и* ± 1.

Рассмотрим половину всех векторов из множества $\mathcal{D}$ среди которых нет противоположных. Данное множество $\mathcal{D}^+$ также однозначно определяет дайсинг $\mathfrak{D}$. Рассмотрим квадратичную форму

$$\varphi(\mathbf{x}) = \sum_{\mathbf{d}\in\mathcal{D}^+} \omega_{\mathbf{d}}(\mathbf{d}\cdot\mathbf{x})^2$$

для некоторых положительных $\omega_{\mathbf{d}}$. Тогда соответствующий многогранник Дирихле-Вороного описывается так [2, Теорема 4.3]:

Теорема 3.8 (Р. Эрдал)**.** *Многогранником Дирихле-Вороного решетки* $\Lambda(\mathfrak{D})$ *относительно квадратичной формы* $\varphi(\mathbf{x})$ *является зонотоп с векторами зон*

$$\mathbf{z}_{\mathbf{d}} = \left(\sum_{\mathbf{d}\in\mathcal{D}^+} \omega_{\mathbf{d}}\mathbf{d}\mathbf{d}^T\right)^{-1} \omega_{\mathbf{d}}\mathbf{d}, \quad \mathbf{d}\in\mathcal{D}^+.$$

Более того, в той же работе доказано, что если метрика $\varphi(\mathbf{x})$ — стандартная евклидова, то векторы зон равны $\mathbf{z}_{\mathbf{d}} = \omega_{\mathbf{d}}\mathbf{d}$ ([2, Разд. 6, стр. 442]).

Также в [2, Теорема 1.2] доказана, что многогранник Дирихле-Вороного для решетки является зонотопом если и только если соответствующее разбиение Делоне является дайсингом.

4 Вторая гипотеза Вороного для зонотопов

Лемма 4.1. *Пусть* $Z = Z(V)$ *—* d*-мерный зонотоп. Любая гипергрань* Z *порождается* $(d-1)$*-мерным подмножеством* U *множества* V *и наоборот, любой* $(d-1)$*-мерное подмножество* V *порождает гипергрань* $Z(U)$ *зонотопа* $Z(V)$*.*

Доказательство. Пусть π — гиперплоскость какой-либо гиперграни Z. Зонотоп $Z(V) = Z(\mathbf{v}_1, \dots \mathbf{v}_n)$ является проекцией куба $C^n \subset \mathbb{R}^n$ на пространство $\mathbb{R}^d$ вдоль $(n-d)$-мерного пространства ψ. Рассмотрим в пространстве $\mathbb{R}^n$ гиперплоскость $\pi\times\psi$; она является опорной гиперплоскостью для куба C^n, а значит определяет его грань F. Грань F является кубом размерности $d-1$, который порождается сторонами куба C^n, проецируемыми в векторы множества V, параллельные плоскости π. Следовательно, вся грань F проецируется в параллельную копию зонотопа $Z(U)$ и в тоже время в грань зонотопа $Z(V)$. В обратную сторону доказательство аналогично. ☐

Теорема 4.2. *Для зонотопов, являющихся параллелоэдрами верна гипотеза 2.*

Доказательство. Пусть зонотоп Z является d-мерным параллелоэдром. Для него верна первая гипотеза Вороного [2, Теорема 1.1], то есть существует такое аффинное преобразование $\mathcal{A}$, что зонотоп $\mathcal{A}Z$ является многогранником Дирихле-Вороного для некоторой решетки Λ. Разбиение Делоне для решетки Λ и стандартной метрики является дайсингом $\mathfrak{D} = \mathcal{D}(\pm\mathbf{d}_1, \dots, \pm\mathbf{d}_n)$, следовательно (теорема 3.8), векторы зон зонотопа $\mathcal{A}Z$ это векторы вида $\omega_i\mathbf{d}_i$.

Вектор, соединяющий центры O_1 и O_2 двух копий $\mathcal{A}Z$ с общей гипергранью F, перпендикулярен это гиперграни, так как все точки F равноудалены от O_1 и O_2. По лемме 4.1 грани F параллельны $d-1$ вектор из множества $\mathcal{D}$, а значит по теореме 3.5, найдется вектор из $\mathcal{E}$ перпендикулярный F, следовательно вектор $\overrightarrow{O_1O_2}$ лежит в $\mathcal{E}$. Обратное также верно, если мы возьмем любой вектор из $\mathcal{E}$, то $d-1$ независимых векторов из $\mathcal{D}$, которые ему перпендикулярны, зададут гипергрань зонотопа Z. Таким образом, множества $\mathcal{E}$ и $\mathcal{N}(Z)$ совпадают.

Рассмотрим унимодулярное представление соответствующего дайсинга. В данном представлении в множестве $\mathcal{E}$ содержится стандартный базис пространства, а остальные векторы множества $\mathcal{E}$ имеют целочисленные координаты. Следовательно, в унимодулярном представлении и в множестве $\mathcal{N}(Z)$ содержится стандартный базис пространства, а остальные векторы этого множества выражаются через данный базис с целыми коэффициентами. □

Список литературы

[1] M. Deza, V. Grishukhin, *Once more about Voronoi's conjecture and space tiling zonotopes.* e-print, http://arxiv.org/abs/math/0203124, 2002.

[2] R. Erdahl, *Zonotopes, Dicings, and Voronoi's Conjecture on Parallelohedra*, Eur. J. of Comb., Vol. 20, N. 6, 1999, pp. 527-549.

[3] P.McMullen, *Convex bodies which tile space by translation.* Mathematika, vol. 27, n. 1, 1980, pp. 113-121.

[4] H. Minkowski, *Allgemeine Lehrsätze über die convexen Polyeder.* Gött. Nachr., 1897, pp. 198-219.

[5] G. Voronoi, *Nouvelles applications des paramètres continus à la théorie des formes quadratiques. Deuxième mémoire. Recherches sur les paralléloèdres primitifs.* J. für Math., vol. 136, 1909, pp. 67-178.

[6] Б.А. Венков, *Об одном классе эвклидовых многогранников.* Вестник Ленинградского Университета, сер. мат., физ., хим., 1954. Том 9, 11-31.